



Crop	Application Timing	Rate
Legumes (Snapbeans, Green Beans, all Peas)	<ol style="list-style-type: none"> 1. At 1st and 2nd trifoliolate leaf stage 2. At 1st Bloom (R1) 	2 to 2 1/2 pints per acre
Tomatoes, Peppers, and Eggplant	<ol style="list-style-type: none"> 1. At 21 days after sowing (6-8 inch stage or 2-4 true leaf stage) 2. 10-14 days later 3. 10-14 days later until flower set 	2 to 2 1/2 pints per acre
Watermelon, Pumpkin, Cucumber, Squash	<ol style="list-style-type: none"> 1. At 21 days after sowing (6-8 inch stage or 2-4 true leaf stage) 2. 10-14 days later 3. 10-14 days later until flower set 	3 pints per acre
Onion	<ol style="list-style-type: none"> 1. At 2-4 true leaf stage or 2-3 weeks after emergence 2. 14-21 days later 3. 14-21 days later 	2 to 2 1/2 pints per acre
Cabbage, Broccoli Cauliflower	<ol style="list-style-type: none"> 1. At 4 to 6 true leaf stage 2. 10-14 days later 3. At head initiation 	2 to 2 1/2 pints per acre
Potatoes	<ol style="list-style-type: none"> 1. 3-4 weeks after emergence or a tube set 2. 10-14 days later 	2 to 2 1/2 pints per acre
Corn	<ol style="list-style-type: none"> 1. At V2-3 (2-4 leaf stage) 2. At VT-R1 	2 to 2 1/2 pints per acre
Strawberries	<ol style="list-style-type: none"> 1. 10-14 days after emergence or 21 days after plant out 2. Every 14-21 days until harvest 	2 to 2 1/2 pints per acre
Celery	<ol style="list-style-type: none"> 1. 14-21 days after emergence 2. 10-14 days later 3. 10-14 days later 	2 to 2 1/2 pints per acre